

RESPIRATORY FUNCTION LABORATORY

Medical Director
Dr Donald Lee
BSc(Med) MBBS(Hons) MMed FRACP

PATIENT DETAILS

Name:
Address:
Date of Birth:

Smoking: Current Former Never Pack years?

Hb, date taken:

CLINICAL HISTORY

TESTS REQUIRED: please tick box

- Full pulmonary function (flow volume loop, with bronchodilator, lung volumes, gas transfer capacity)
- Flow volume loop/ Spirometry, includes bronchodilator
- MIPS & MEPS (mouth pressures)
- Six Minute Walk Test
- Home Overnight Oximetry

Tests for Asthma/ Allergy

- Bronchoprovocation Challenge (Mannitol)*
- Nasal Resistance
- Skin Test for Allergies**
- Exhaled Nitric Oxide

Referring Doctor:

Provider number:

Address:

Date: / /

Fax:

*Fees apply
**Must provide clinical history

Pulmonary Function, Spirometry & Flow Volume Loops

- No short acting bronchodilators FOUR HOURS prior to test such as Ventolin, Bricanyl and Atrovent.
- No long acting bronchodilators 12hrs prior to test such as Serevent, Seretide, Oxis and Symbicort.
- No Onbrez, Spiriva, Seebri, Bretaris, Ultibro, Breo, Anoro, Incruse or Singulair for 48hrs prior to testing.
- No smoking for two hours prior to testing (ideally 24 hours).

Mannitol Challenge

- No short acting bronchodilators 12 hours prior to test such as Ventolin, Bricanyl and Atrovent.
 - No Serevent, Oxis, Spiriva, Seebri, Bretaris, Ultibro, Breo, Anoro or Incruse for at least 48 hours prior to testing.
 - For asthma diagnosis ideally no Flixotide, Intal, Pulmicort, Seretide, Symbicort, Qvar, Alvesco or Tilade for two weeks before test.
(Although tests performed on therapy to assess effectiveness).
 - No antihistamines, theophylline or Singulair tablets for at least 48hrs prior to testing.
 - Oral Prednisolone, etc (cortisone)
- (NB: Please discuss with prescribing doctor before withholding oral cortisone medication- Test not valid if taking this medication)

Nasal Resistance

- No vasoconstrictors 4hrs prior to test such a Drixine, Dimetapp.

Skin Tests

- Ideally should not be done if patient is on Prednisone $\geq 5\text{mg/day}$. No antihistamines for at least 48hrs prior to testing.

Limited parking available onsite

